

Web of Science

Search

Search Results

My Tools ▾

Search History

Marked List



Save to EndNote online ▾

Add to Marked List

13 of 22

Cloning and Over-expression of xynB Gene of Bacillus subtilis subsp spizizenii W23 into Escherichia coli Origami Host Cells

By: [Wahjudi, M](#) (Wahjudi, Mariana)^[1]; [Catherina](#) (Catherina)^[1]; [Wangunhardjo, NM](#) (Wangunhardjo, Nita Marcelia)^[1]; [Suryadjaja, E](#) (Suryadjaja, Ernest)^[1]; [Daniel, X](#) (Daniel, Xavier)^[2]

Edited by: [Setyobudi, RH](#); [Purwanto, MGM](#); [Burlakovs, J](#); [Mel, M](#); [Adinurani, PG](#); [VincevicaGaile, Z](#)

INTERNATIONAL CONFERENCE ON NATURAL RESOURCES AND LIFE SCIENCE (NRLS-2016)

Book Series: KnE Life Sciences

Pages: 139-149

DOI: 10.18502/kls.v3i5.986

Published: 2017

Conference

Conference: 1st International Conference on Natural Resources and Life Sciences (NRLS)

Location: Surabaya, INDONESIA

Date: OCT 20-21, 2016

Sponsor(s): Deutschland Akademische Austausch Dienst; Univ Surabaya, Fac Biotechnol; SCIENCEWERKE; Bioneer; Indolab Utama; Enseval Medika Prima; Megah Sejahtera Sci; Alumni Portal Deutschland

Abstract

The xynB gene of Bacillus subtilis subsp. spizizenii W23 is predicted to encode a xylan 1,4-beta-xylosidase. Application of XynB enzymes in industries is wide. Production of this enzyme in its host cells is naturally restricted by repression process. It will give certain beneficial to over-expressed the enzymes in other host-cells under inducing promoter. This study aimed to clone the xynB gene from Bacillus subtilis subsp. spizizenii W23, to pMMB67EH plasmid, and to over-express the xynB gene in Escherichia coli Origami as host cells. The xynB gene was successfully amplified by polymerase chain reaction (PCR) technique using a pair of primers flanking the gene sequence and chromosomal DNA of the W23 strain as a template. The xynB gene inserted in recombinant plasmid was confirmed by PCR detection using primers pair's specific for xynB gene and for the vector, then continued by restriction analyses. The result showed that transformants clone 9 and 10 bear the recombinant pMMB-xynB plasmid. The xylanase activity of xynB gene in Escherichia coli Origami clone 10 was detected by sodium-dodecyl-sulfate polyacrylamide gel analyses and with addition of isopropyl-beta-D-thiogalactoside (IPTG) as an inducer. The protein seem to be over-expressed as intra-and extra-cellular protein detected on SDS-PAGE gel. Result from xylan degrading activity on Luria-Bertani-xylan-IPTG plate with addition of Congo Red, showed that the cells with pMMB-xynB recombinant plasmid have clear zone around the colonies while the transformant bearing an empty plasmid showed no clear zone. It could be concluded that the xynB gene of Bacillus subtilis subsp. spizizenii W23 has been successfully been cloned on pMMB67EH plasmid and over-expressed in the Escherichia coli Origami

Citation Network

In Web of Science Core Collection

0

Times Cited

Create Citation Alert

40

Cited References

[View Related Records](#)

Use in Web of Science

Web of Science Usage Count

0

Last 180 Days

0

Since 2013

[Learn more](#)

This record is from:

Web of Science Core Collection

- Conference Proceedings Citation Index-Science

Suggest a correction

If you would like to improve the quality of the data in this record, please [suggest a correction](#).

Keywords

Author Keywords: Bacillus subtilis subsp spizizenii str. W23; cloning; Escherichia coli Origami; over-expression; xylan-beta 1,4-xylosidase (xynB)

KeyWords Plus: XYLAN-DEGRADING ENZYMES; CATABOLITE REPRESSION; MICROBIAL XYLANASES; PREDICTION; ETHANOL; OPERON; GROWTH; PULP

Author Information

Reprint Address: Wahjudi, M (reprint author)

Univ Surabaya Ubaya, Fac Biotechnol, Surabaya 60293, Indonesia.

Addresses:

[1] Univ Surabaya Ubaya, Fac Biotechnol, Surabaya 60293, Indonesia

[2] ITS Educ Asia, Unit G, 4-F BOC Grp Life Assurance Tower,134-136 Voeux Rd, Hong Kong
999077, Hong Kong, Peoples R China

E-mail Addresses: mariana_wahyudi@staff.ubaya.ac.id

Funding

Funding Agency	Grant Number
Ubayu Research and Community Development Center (University of Surabaya)	027/Lit/LPPM-01/FT/XII/2012

[View funding text](#)

Publisher

KNOWLEDGE E, OFFICE 4402, X2 TOWER, JLT, PO BOX 488239, DUBAI, 00000, U ARAB
EMIRATES

Categories / Classification

Research Areas: Life Sciences & Biomedicine - Other Topics

Web of Science Categories: Biology

Document Information

Document Type: Proceedings Paper

Language: English

Accession Number: WOS:000417445000013

ISSN: 2413-0877

Other Information

IDS Number: BJ1EF

Cited References in Web of Science Core Collection: [40](#)

Times Cited in Web of Science Core Collection: [0](#)

Scientific and Editorial (S & E) Board International Conference on Natural Resources and Life Sciences the 1st NRLS - 2016

- Elieser Tarigan, Faculty of Electrical Engineering - University of Surabaya, IDN.
- Go Melisa Gunawan, Faculty of Biotechnology - University of Surabaya, IDN.
- Johan Sukweenadhi, Kyung Hee University - KOR; Faculty of Biotechnology - University of Surabaya, IDN.
- Juris Burlakovs, Faculty of Health and Life Sciences, Dept. of Biology and Environmental Science, Linnaeus University, SWD
- Lanny Sapei, Faculty of Chemical Engineering - University of Surabaya, IDN.
- Lieke Riadi, Faculty of Chemical Engineering - University of Surabaya, IDN.
- Maizirwan Mel, Department of Biotechnology Engineering, Faculty of Engineering, International Islamic University Malaysia, MYS.
- Maria Goretti Marianti Purwanto, Faculty of Biotechnology - University of Surabaya, IDN.
- Mariana Wahjudi, Faculty of Biotechnology - University of Surabaya, IDN.
- Nurul Azizah, Faculty of Biotechnology - University of Surabaya, IDN.
- Popy Hartatie Hardjo, Faculty of Biotechnology - University of Surabaya, IDN.
- Praptiningsih Gamawati Adinurani, Faculty of Agrotechnology, University of Merdeka, Madiun, IDN.
- Roy Hendroko Setyobudi, IABST, Jakarta and Ma Chung Research Center for Photosynthetic Pigments, Malang, IDN.
- Sharr Azni Harmin, Fakulti Sains & Bioteknologi, Universiti Selangor, MYS.
- Sulistyo Emantoko Dwi Putra, Faculty of Science - University of Potsdam, DEU; Faculty of Biotechnology University of Surabaya, IDN.
- Theresia Desy Askitosari, RWTH Aachen University, Institute of Applied Microbiology, DEU.
- Tjie Kok, Faculty of Chemical Pharmaceutical Biology, University of Groningen, NLD.



- Yoanes Maria Vianney, Faculty of Biotechnology - University of Surabaya, IDN.
- Zane Vincēviča-Gaile, Department of Environmental Science, University of Latvia, Riga - Latvia, European Union, LVA.

Table of Contents (NRLS Conference Proceedings)

Action Research into a Flood Resilient Value Chain – Biochar-Based Organic Fertilizer
Doubles Productivity of Pea in Udayapur, Nepal

Anu Joshi Shrestha, Bishnu Hari Pandit

DOI 10.18502/cls.v3i5.974, Pages 1-19

Comparison of the Main Bioactive Compounds and Antioxidant Activity from Garlic Water-soluble and Garlic Oil

Ardhia Deasy Rosita Dewi, Joni Kusnadi, Wen-Ling Shih

DOI 10.18502/cls.v3i5.975, Pages 20-34

Antioxidant Activity in Various Processed Products of Inferior Local Tubers (*Dioscorea* sp. L.)

Ari Yuniastuti, Retno Sri Iswari, and R. Susanti

DOI 10.18502/cls.v3i5.976, Pages 35-40

Synthesis and Characterization of TNTs/Polyaniline Composite as Photocatalyst Degradation of Rhodamin B by Visible Light

Atik Setyani, Sri Wahyuni, Sigit Priatmoko, and Nur Amin

DOI 10.18502/cls.v3i5.977, Pages 41-50

Two Antibacterial Compounds: Velutin and 4- (Hydroxy (Oxiran-2-yl)methyl)-2-Methoxyphenol from the Stem Bark of *Drimys arfakensis* Gibbs.

Bimo Budi Santoso, Hidelisa Padua Hernandez, Evelyn B Rodriguez, Ida Farida Dalmacio

DOI 10.18502/cls.v3i5.978, Pages 51-62

Extraction and Toxicity Assay of Mycotoxin from Entomopathogenic Fungi Isolate of Kusuma Agrowisata Orchard Batu, Jawa Timur, Indonesia

Derdy Janli, Maria Goretti M. Purwanto, Ida Bagus Artadana, Theresia Desy Askitosari

DOI 10.18502/cls.v3i5.979, Pages 63-74

Solar Driven Photocatalytic Fuel Cells: an Environmentally Friendly Innovation of Power Plant Technology as Energy Source

Emas Agus Prastyo Wibowo, Navela Rahma Aji, Nuni Widiarti, Cepi Kurniawan

DOI 10.18502/cls.v3i5.980, Pages 75-80

A Glimpse into the Biosynthesis of Terpenoids

Ingy I. Abdallah, Wim J. Quax

DOI 10.18502/cls.v3i5.981, Pages 81-98

Transcription Pattern of Catalase Gene from *Gynostemma pentaphyllum* (Thunb.) Makino during Various Abiotic Stresses

Johan Sukweenadhi, Yu-Jin Kim, and Deok-Chun Yang

DOI 10.18502/cls.v3i5.982, Pages 99-109

Drying Kinetics of *Curcuma xanthorrhiza* Roxb.

Lanny Sapei, Elieser Taringan, Lie Hwa, Violita Putri Halim, Fhelix August Soebiantoro

DOI 10.18502/cls.v3i5.983, Pages 110-117

Water Sustainability: Emerging Trends for Water Quality Management

Lieke Riadi

DOI 10.18502/cls.v3i5.984, Pages 118-130

Isolation and Screening of Indigenous Fungus Producing Lignin Peroxidase from the Cocoa Plantation in Sepawon Kediri Regency Indonesia

Liling Delila, Evi Susanti, Eli Hendrik Sanjaya

DOI 10.18502/cls.v3i5.985, Pages 131-138

Cloning and Over-expression of xynB Gene of *Bacillus subtilis* subsp. *spizizenii* W23 into *Escherichia coli* Origami Host Cells

Mariana Wahjudi, Catherina, Nita Marcelia Wangunhardjo, Ernest Suryadjaja, Xavier Daniel

DOI 10.18502/cls.v3i5.986, Pages 139-149

The Utilization of Metroxylon Sago (Rottb.) Dregs for Low Bioethanol as Fuel Households Needs in Papua Province Indonesia

Numberi Johni Jonatan, Arifia Ekayuliana, I Made Kartika Dhiputra, Yulianto Sulisty Nugroho

DOI 10.18502/cls.v3i5.987, Pages 150-157

Composting of Corn By-Product using EM4 and Microorganism *Azotobacter* sp. as Composting Organism

Nuniek Hendriani, Sri Rachmania Juliastuti, Hamida Nuur Masetya, Imam Tianto Aditias Saputra

DOI 10.18502/cls.v3i5.988, Pages 158-166

Superoxide Dismutase Levels of Operator Gas Stations in Semarang, Central Java, Indonesia.

Nur Kusuma Dewi, Ari Yuniastuti

DOI 10.18502/cls.v3i5.989, Pages 167-172

Somatic Embryo from Basal Leaf Segments of *Vanda tricolor* Lindl. var. *pallida*

Popy Hartatie Hardjo and Wina Dian Savitri

DOI 10.18502/cls.v3i5.990, Pages 173-179

Efficiency Analysis of Production Factors Utilization in Upland Rice Farming

Rohmad Budiono, Praptiningsih Gamawati Adinurani

DOI 10.18502/cls.v3i5.991, Pages 180-187

In Vitro Study of Potato (*Solanum tuberosum* L.) Tolerant to the Drought Stress

Biliter Anton Sirait and Rosa Charloq

DOI 10.18502/cls.v3i5.992, Pages 188-192

Organic Fertilizer from Bioethanol Solid Waste, Agricultural Waste, and Banana Peels Waste
by Bio-act EM4 and *Aspergillus niger*
Sri Rachmania Juliastuti, Delftya Enhaperdhani, Rizka Uswatun Hasanah
DOI 10.18502/kl.v3i5.993, Pages 193-201

Intake Pericarp of *Garcinia mangostana* L. Extract Inhibited Oxidative Stress on Wistar Rat
Hyperglycemic through the Increased of Superoxide Dismutase and Histopathology
Pancreatic Activities
Sri Wahjuni, Anak Agung Istri Ayu Mayun Laksmiwati, Ni Made Puspawati
DOI 10.18502/kl.v3i5.994, Pages 202-207

The Role of Heavy Metals-Resistant Bacteria *Acinetobacter* sp. in Copper Phytoremediation
using *Eichhornia crasippes* [(Mart.) Solms]
Wahyu Irawati, Adolf Jan Nexson Parhusip, Nida Sopiah, Juniche Anggelique Tnunay
DOI 10.18502/kl.v3i5.995, Pages 208-220

Conference Paper

Cloning and Over-expression of *xynB* Gene of *Bacillus subtilis* subsp. *spizizenii* W23 into *Escherichia coli* Origami Host Cells

Mariana Wahjudi¹, Catherina¹, Nita Marcelia Wangunhardjo¹, Ernest Suryadjaja¹, and Xavier Daniel²

¹Faculty of Biotechnology, University of Surabaya (Ubaya), Jl. Raya Kalirungkut, Surabaya 60293, Indonesia

²ITS Education Asia, Unit G, 4/F BOC Group Life Assurance Tower, 134-136 Des Voeux Road, Central, Hong Kong 999077

Abstract

The *xynB* gene of *Bacillus subtilis* subsp. *spizizenii* W23 is predicted to encode a xylan 1,4-beta-xylosidase. Application of XynB enzymes in industries is wide. Production of this enzyme in its host cells is naturally restricted by repression process. It will give certain beneficial to over-expressed the enzymes in other host-cells under inducing promoter. This study aimed to clone the *xynB* gene from *Bacillus subtilis* subsp. *spizizenii* W23, to pMMB67EH plasmid, and to over-express the *xynB* gene in *Escherichia coli* Origami as host cells. The *xynB* gene was successfully amplified by polymerase chain reaction (PCR) technique using a pair of primers flanking the gene sequence and chromosomal DNA of the W23 strain as a template. The *xynB* gene inserted in recombinant plasmid was confirmed by PCR detection using primers pair's specific for *xynB* gene and for the vector, then continued by restriction analyses. The result showed that transformants clone 9 and 10 bear the recombinant pMMB-*xynB* plasmid. The xylanase activity of *xynB* gene in *Escherichia coli* Origami clone 10 was detected by sodium-dodecyl-sulfate polyacrylamide gel analyses and with addition of isopropyl- β -D-thio-galactoside (IPTG) as an inducer. The protein seem to be over-expressed as intra- and extra-cellular protein detected on SDS-PAGE gel. Result from xylan degrading activity on Luria-Bertani-xylan-IPTG plate with addition of Congo Red, showed that the cells with pMMB-*xynB* recombinant plasmid have clear zone around the colonies while the transformant bearing an empty plasmid showed no clear zone. It could be concluded that the *xynB* gene of *Bacillus subtilis* subsp. *spizizenii* W23 has been successfully been cloned on pMMB67EH plasmid and over-expressed in the *Escherichia coli* Origami cells as intra- and extra-cellular protein, as observed on SDS-PAGE gel analysis. The protein has activity on xylan degradation.

Keywords: *Bacillus subtilis* subsp. *spizizenii* str. W23; cloning; *Escherichia coli* Origami; over-expression, xylan-beta 1,4-xylosidase (*xynB*).

Corresponding Author:

Mariana Wahjudi
mariana_wahjudi@staff.
ubaya.ac.id

Received: 9 June 2017

Accepted: 15 July 2017

Published: 11 September 2017

Publishing services provided
by Knowledge E

© Mariana Wahjudi et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Selection and Peer-review under the responsibility of the NRLS Conference Committee.

OPEN ACCESS

1. Introduction

Xylanases are important enzymes for degradation of xylans [1], the major components of hemicellulosic plant materials. Xylan consists of a backbone of polymer β -1,4-linked-xylose with side chains of xylose, arabinose, acetyl, methylglucuronic, coumaric, hexoses, and uronic acid. Hemicellulose together with cellulose and lignin can be found in plant cell wall in the form of lignocelluloses. Complete degradation of xylans requires the involvement of several enzymes [2]. Xylan 1,4- β -xylosidase (4- β -D-xylan xylohydrolase, EC 3.2.1.37) remove D-xylose residues from the non-reducing termini, and endoxylanases (4- β -D-xylan xylanohydrolase; EC 3.2.1.8) catalyze endohydrolysis of (1 \rightarrow 4)- β -D-xylosidic linkages in xylans. All those enzymes degrade the xylan backbone into xyloses whereas others enzymes such as acetyl esterase, ferulic esterase, glucuronosidase or arabinosidase remove the side chains residues [3].

Xylanases are important enzymes for many industries. In pulp and paper industries [4, 5], candy, and syrup industries [6]. Xylo-oligosacharides products have wide applications in industries, such as in juice-processing and textile industries, bread industry [7]. In poultry industries, they are added in feed processing to improve the feed digestions and nutrients uptake [3, 8, 9].

Xylanases have been found in various organisms such as *Thermatoga maritima* [10], *Aspergillus niger* [11], *Clostridium cellulovorans* [12], *Bacillus* sp.CCMI 966 [13], *Cellulomonas uda* [14], *Thermoanaerobacterium* sp. strain [15], *Dictyoglomus thermophilum* [16], *Ustilago maydis* [17] and many others bacteria and fungi [6]. The enzymes properties may be slight different depend on the sources organisms.

Over-expression of genes encoding celluloses and xylanases has many advantages since lignocelluloses are present abundantly in forest and agricultural as wastes. In other hands, these wastes still contain nutrients for fermentation process to produce important stuffs [18, 19]. The lignocelluloses degradation may help the carbon recycle in the earth. Furthermore, the degradation products may be used as nutrients for microorganisms in fermentation processes. Many studies showed that hydrolytic products could increase cell biomass and bioproducts [19, 20] with which no competition for food and feed requirements. Expressions of *xyn* gene in microorganism producers enhanced the xylan degradation and increase the bioproducts, such as ethanol [21].

In order to produce this enzyme in large scale, we intended to clone the gene from *Bacillus subtilis* subsp. *spizizenii* W23 and introduced into others host cells, such as *Escherichia coli* Origami. Productions of proteins in original cells have advantages over other bacterial cells since *Bacillus* has a protein secretory system. However, native cells commonly have inner repression process which causes restriction in practice. Several study revealed the presence of *xyn* operon regulation [22], for instance carbon catabolite repression of *Bacillus subtilis* xylanases [23, 24]. Because of that repression,

it will be more advantageous if this *xynB* gene can be over-expressed not in its native host cells. Further study can be performed to regulate or manipulate the process even in native host cells.

Commonly, xylanases genes were found in genomic library fragments of microorganisms found in their nature habitats, such as done for *Paenibacillus* sp. [25], *Bacillus pumilus* [26]. However, the fragments often contain the whole operon with intact regulator in the fragment [26]. In this study, we isolated the *xynB* gene by PCR amplification. *Bacillus subtilis* subsp. spizizenii W23 genome have complete genome sequence for this purpose. The difficulty of this step is in the designing of primers for amplification the whole coding sequence. Sequences around *xynB* gene region are rather complicated for primers designing [27]. In this flanking area, no sequences are available for proper annealed primer. Based on several software analyses, the sequences contain repeated nucleotides, low GC content and several unspecific sequences (NCBI Primer Design Tool [28], Primer 3 [29, 30], Clone Manager Suite Professional softwares). In this study, we managed to design the primers pair for whole coding sequence of *xynB* gene and amplified it for construction of recombinant plasmid.

The aims of this study are to construct a recombinant plasmid containing *xynB* gene from *Bacillus subtilis* subspecies spizizenii W23 and to over-express the *xynB* gene in *Escherichia coli* Origami as host cells under tac promoter.

2. Materials and methods

2.1. Bacteria and plasmid

Escherichia coli Origami and *Escherichia coli* DH5 α (Novagen) containing pMMB67EH plasmid [31]. *Bacillus subtilis* subsp.spizizenii W23 is a kindly gift from Northern Regional Research Laboratory, NRRL)

2.2. Amplification of *Bacillus subtilis* subsp.spizizenii W23 *xynB* gene

The open reading frame (ORF) of *Bacillus subtilis* subsp.spizizenii W23 gene was amplified by polymerase chain reaction (PCR) using primers that were designed based on the genomic sequence of *Bacillus subtilis* subsp.spizizenii W23 strain obtained from Genebank database (www.ncbi.com) [27], using Clone Manager Professional Suite (DNA Star, USA) software. For-*xynB*_BSUW23: 5'-CGAGCGGCCGAGCTCATGAAGATTATCAATCCAGTACTTAAAGGATTCAA-3', and Rev-*xynB*_BSUW23: 5'-CCGCTGGTCTAGAGGCTCCTGTAAAGAACCTCCTCCC-3'. Both primers were added with a restriction sites at its

5'-ends (underlined), and synthesized by Operon. As a PCR template was chromosomal DNA of *Bacillus subtilis* subsp. *spizizenii* W23. PCR was settled as follow, pre-denaturation at 94 °C for 2 min, denaturation at 94 °C for 30 s, annealing at 60 °C for 1.5 min, elongation at 72 °C for 2.5 min, 30 cycles and post-extension at 72 °C for 10 min.

2.3. Cloning of PCR fragment on pMMB67EH vector

The *xynB* gene was cloned into pMMB67EH under the control of tac promoter [31]. Before ligation, the PCR product was digested with *SacI* and *XbaI* enzymes and then was ligated to the similarly digested pMMB67EH. The resulting pMMB-*xynB* plasmid was used to transform the *Escherichia coli* Origami cells which were made competence by CaCl_2 using heat shock method, as described by Sambrook et al. [32].

2.4. Analyses of recombinant plasmid

Screening of transformants was done by plating the transformants on LB plate containing $100 \mu\text{g} \cdot \text{mL}^{-1}$ ampicillin (LB-amp). The transformants grew on LB-amp plates were then selected for the presence of recombinant plasmid in the cells by PCR techniques. For vector detection, PCR was performed using vector-specific primers, SeqForpMMB67 dan SeqRevpmMB67 primers [33]. The amplification product was predicted to be a 2 035 bps band, of which the 1 523 bps was the size of *xynB* gene sequence and the rest was the vector parts. While for *xynB* detection, the For-*xynB*_BSUW23 and Rev-*xynB*_BSUW23 primers pair was used. The recombinant plasmids generate a single band amplicon with a size of ~1 549 bps. As a control, *Escherichia coli* Origami, with or without plasmids, were used as templates. All of these negative controls have no amplicon.

The restriction analyses were done for recombinant plasmid that gave positive result in PCR detections. The restriction analyses were performed using *SacI*, *XbaI* and *EcoRI* (Fermentas) restriction enzymes according to the manufacturer's instruction and then visualized on 1 % agarose gel. *XbaI* or *SacI* will linearized pMMB-*xynB* to be a linear fragment with a size of 10 334 bps. *EcoRI* will digest proper recombinant plasmid to two fragments of ~1005 and ~9 329 bps, while digestion with *XbaI* and *SacI* simultaneously gave two fragments, 1 523 bps which confirmed for the size of *xynB* insert and 8 811 bps for the size of pMMB-vector. Un-cut empty plasmid, pMMB67EH, had a size of ~8 828 bp.

2.5. Over-expression verification by SDS-PAGE analysis

Over-expression of *xynB* gene in *Escherichia coli* Origami cells was verified on sodium-dodecyl-sulfate polyacrylamide (SDS-PAGE) gel analyses. The cells were grown in LB-amp, at 37 °C, and induce with 1 mM IPTG. The pellet was separated from the supernatant by centrifugation at 13 000 *g* for 1 min [g = Relative Centrifugal Force (RCF)], then sonicated to break the cells membrane. The separation of proteins was performed on SDS-PAGE gel with 3 % stacking gel and 12.5 % separating gel [32]. The proteins bands were visualized by Coomassie blue G-250 staining.

2.6. Xylan degradation activity

The xylan degradation activity was assayed by growing the transformants on LB-amp plate containing 1 % xylan with addition of 1mM IPTG, incubated at 37 °C for 7 d. The *Escherichia coli* Origami transformants harboring xylanase gene showed a clear zone around the colony upon Congo red addition [34]

3. Results and discussion

The *xynB* gene had been amplified successfully using For-*xynB*_BSUW23 and Rev-*xynB*_BSUW23 primers. PCR product revealed as a single band with a size of 1 549 bps (Fig. 1). Commonly, isolation of genes encoding xylanases from various bacteria and fungi have been done by digestion of genomic DNA, cloning the fragments and then the target were selected based on its activity [26]. Both primers were predicted as not good primers by the Clone Manager Suite Professional software, however in fact the amplification was succeeded in this study.

The screening of *Escherichia coli* Origami transformants were performed based on ampicillin resistance [33]. There were many transformants colonies grew on LB-amp plates, as many as colonies of *Escherichia coli* Origami transformed using the empty plasmid, pMMB67EH. The pMMB67EH is a vector plasmid contained ampicillin resistance marker. Since ampicillin resistance is quite common in bacterial cells and laboratory plasmids, so further analyses have to be performed.

Ten of *Escherichia coli* Origami transformants grew on LB-amp plates were selected to detect the presence of target plasmid by PCR using *SeqForpMMB67* and *SeqRev-pMMB67* primers. The result showed that nine of *Escherichia coli* Origami transformants had a fragment with proper size of ~2.0 kbs on 1 % agarose gel. Based on this analysis, these nine *Escherichia coli* Origami transformants were confirmed to contain the pMMB-recombinant plasmid.




Figure 1: The *xynB* gene was amplified from *Bacillus subtilis* subsp. *spizizenii* W23 genome. The PCR product had a band of ~1 549 bp (line 2), while digestion with *SacI* and *XbaI* resulted to three fragments with sizes of approximately 1 523 bps, 14 bps and 12 bps (line 1). The 14 and 12 bps fragments were too small to be existed on this gel under the electrophoresis condition of this experiment. The PCR product and restriction fragments had proper sizes as predicted. M1 and M2: 1 kb and 100 bps DNA ladder (Fermentas).

To confirm the presence of *xynB* in the recombinant plasmids, two of the positive transformants, *Escherichia coli* Origami clones 9 and 10, were chosen for next PCR detection using For-*xynB*_BSUW23 and Rev-*xynB*_BSUW23 primers. Before PCR process, the plasmids were isolated from clones 9 and 10 which then were used as templates in PCR amplification. PCR result showed that both transformants have pMMB-*xynB* plasmids with *xynB* gene in proper orientation.

For further analysis, restriction analysis was done to confirm the orientation of the *xynB* gene in the pMMB67EH vector. The pMMB-*xynB* plasmid was extracted from *Escherichia coli* Origami transformants clone 9 and 10, then were digested using *XbaI*, and/or *SacI*. Restriction analyses results showed that digested recombinants plasmids have proper fragments sizes. All of the analyses results confirmed that the *Escherichia coli* Origami clone 9 and 10 contain the proper pMMB-*xynB* recombinant plasmid.

To verify that the *xynB* gene positioned in frame on the recombinant plasmid so that over-expressed properly in transformant cells, the proteins of clone 10 were profiled on SDS-PAGE gel (Fig. 2).

Deducted from the DNA sequence, the protein is predicted to have a molecular mass of ~61 337 Da. SDS-PAGE result showed that this positive clone revealed a thicker protein band in the size of ~61 kDa, both in intra- and extra-cellular fractions. This result gave impression that this protein can be an intra- or extra-cellular protein.

To confirm the presence of active protein and its localization, assay was also performed by Congo Red assay. The activity to degrade xylan by the extra-cellular enzyme resulted to clear red zone surrounding the colonies upon Congo red addition. Clear zone surrounding colony of transformant gave confirmation that the transformant over-expressed the active XynB enzyme and it might be secreted into the medium. No clear zone appeared around the wild type colony (Fig. 3). It meant that the *xynB*




Figure 2: SDS-PAGE proteins profiles of over-expressed *E. coli* Origami pMMB-*xynB* cells grown in LB-amp at 37 °C, shaken at 200 rpm (1 rpm = 1/60 Hz), under 1 mM IPTG induction.




Figure 3: The expression of pMMB-*xynB* in *Escherichia coli* Origami transformants assayed on LB-amp plate containing 1 % xylan and 1 mM IPTG. The colonies of cells bearing *xynB* recombinant plasmid showed a clear zone (black arrow, →) around the colonies however no clear zone appeared around transformant bearing empty plasmid. 37 °C, for 7 d.

gene was over-expressed from pMMB-*xynB* in this positive transformants. The *Bacillus subtilis* subsp.spizizenii W23 cells, as an original gene source, showed the highest xylan degrading activities. This result was corroborated to genomic sequence database, indicates that *Bacillus subtilis* subsp.spizizenii W23 has several genes which were predicted to encode lignocellulases enzymes, including the *xynA* and *xynB* genes [27].

Based on protein structure analyses using PROPSearch software [35], *xynB* gene of *Bacillus subtilis* subsp. *spizizenii* W23 is predicted to encode a β -xylosidase (1,4- β -D-xylan xylohydrolase; EC 3.2.1.37). Compare to *xynB* from *Bacillus pumilus* [26], the latter is known as intracellular endoxylanases. *XynB* from *Bacillus pumilus* was detected as an intra-cellular protein by software and by experiment, without signal peptide [26]. Different from the previous protein, this *Bacillus subtilis* W23 *XynB* protein is predicted by PrediSi [36] dan SignalP software(www.ExPassy.com), as a non-secreted protein, without signal peptide sequences. Based on CELLO Prediction software, the most possible protein location is in cytoplasm, however there is a half possibility to be an extracellular protein [37]. In contrast, according to MTHMM software [38], this *XynB* protein is predicted to be an extracellular protein although it contains no signal peptide. This result also reveal in ProtComp [39] analyses result. Although predicted ProtComp software as a cytoplasmic protein, a part of the amino acid residues is located in membrane as integral and transmembran protein. This result can be explained by the fact that the *Bacillus subtilis* sequence have only 48 % similarity to *Bacillus pumilus* that the sequence might be unique and have different properties [40]. Other possibility is that this gene might be expressed abundantly in cytoplasm so that secreted out of the cells. It is interesting to find that this gene was secreted by transformants and gave impact on xylan around the colony. However, the localization of *Bacillus subtilis* subsp.*spizizenii* W23 *XynB* protein still need to be verified.

4. Conclusion

It could be concluded that the *xynB* gene of *Bacillus subtilis* subsp.*spizizenii* W23 has been successfully been cloned on pMMB67EH plasmid and over-expressed in the *Escherichia coli* Origami cells both intra- and extra-cellularly, as observed on SDS-PAGE gel analysis. The transformants bearing the pMMB-*xynB* recombinant plasmid had activity to degrade xylan on LB-xylan plate while the transformant bearing an empty plasmid showed no activity.

Acknowledgement

This work was supported by a research grand of Ubaya Research and Community Development Center (University of Surabaya) for Mariana Wahjudi, grant number 027/Lit/LPPM-01/FT/XII/2012

References

- [1] Collins T, Gerday C, Feller G. Xylanases, xylanase families and extremophilic xylanases. *FEMS Microbiology Reviews* 2005;29(1):3-23.
- [2] Snelders J, Dornez E, Broekaert WF, Delcour JA, Courtin CM. Determination of the xylan backbone distribution of arabinoxylan-oligosaccharides. *Bioactive Carbohydrates and Dietary Fibre* 2013;2(1):84-91.
- [3] Beg QK, Kapoor M, Mahajan L, Hoondal GS. Microbial xylanases and their industrial applications: A review. *Applied Microbiology and Biotechnology* 2001;56(3-4):326-338.
- [4] Walia A, Guleria S, Mehta P, Chauhan A, Parkash J. Microbial xylanases and their industrial application in pulp and paper biobleaching: A review. *3 Biotech* 2017;7(1):11. p.11.
- [5] Kumar S, Haq I, Yadav A, Prakash J, Raj A. Immobilization and biochemical properties of purified xylanase from *Bacillus amyloliquefaciens* SK-3 and its application in kraft pulp biobleaching. *Journal of Clinical Microbiology and Biochemical Technology* 2016;2(1):026-034.
- [6] Burlacu A, Cornea CP, Israel-Roming F. Screening of xylanase producing microorganisms. *Research Journal of Agricultural Science* 2016;48(2):8-15.
- [7] Jiang ZQ, Yang SQ, Tan SS, Li LT, Li XT. Characterization of a xylanase from the newly isolated thermophilic *Thermomyces lanuginosus* CAU44 and its application in bread making. *Letters in Applied Microbiology* 2005;41(1):69-76.
- [8] Lei Z, Shao Y, Yin X, Yin D, Y, Yuan J. Combination of xylanase and debranching enzymes specific to Wheat Arabinoxylan improve the growth performance and gut health of broilers . *Journal of Agricultural and Food Chemistry* 2016;64 (24):4932-4942.
- [9] M.J. Vazquez, J.L. Alonso, H. Domnguez, J.C. Parajo. Xylooligosaccharides: manufacture and applications. *Trends Food Science & Technology* 2000;11(11):387-393.
- [10] Chen CC, Adolphson,R, Dean JFD, Eriksson KEL, Adams MWW, WestphelingJ. Release of lignin from kraft pulp by a hyperthermophilic xylanase from *Thermatoga maritima*. *Enzyme and Microbial Technology* 1997;20(1):39-45.
- [11] Frederick MM, Frederick JR, Fratzke AR, Reilly PJ. Purification and characterization of a xylobiose and xylose-producing endo-xylanase from *Aspergillus niger*. *Carbohydr Res* 1981;97(1):87-103.
- [12] Han SO, Cho HY, Yukawa H, Inui M, Doi RH. Regulation of expression of cellulosomes and noncellulosomal (hemi)cellulolytic enzymes in *Clostridium cellulovorans* during growth on different carbon sources. *Joernal of Bacteriology* 2004;186(13):4218-4227.

- [13] Sá-Pereira P, Costa-Ferreira M, Aires-Barros MR. Enzymatic properties of a neutral endo-1,3(4)-beta-xylanase Xyl II from *Bacillus subtilis*. Joernal of Biotechnology 2002;94(3):265-275.
- [14] Rapp P, Wagner F. Production and properties of xylan-degrading enzymes from *Cellulomonas uda*. Applied Environmental Microbiology 1986;51(4):746-752.
- [15] Lorenz WW, Wiegel J. Isolation, analysis, and expression of two genes from *Thermoanaerobacterium* sp. strain JW/SL YS485: A beta-xylosidase and a novel acetyl xylan esterase with cephalosporin C deacetylase activity. Joernal of Bacteriology 1997;179(17):5436-5441.
- [16] Zhang W, Lou K, Li G. Expression and characterization of the *Dictyoglomus thermophilum* Rt46B.1 xylanase gene (*xynB*) in *Bacillus subtilis*. Applied Biochemistry and Biotechnology 2010;160(5):1484-1495.
- [17] Geiser E, Wierckx N, Zimmermann M, Blank LM. Identification of an endo-1,4-beta-xylanase of *Ustilago maydis*. BMC Biotechnol. 2013;13:59.
- [18] Peterson JD1, Ingram LO. Anaerobic respiration in engineered *Escherichia coli* with an internal electron acceptor to produce fuel ethanol. Annals of the New York Academy of Sciences 2008;1125(1):363-372.
- [19] Mtui GYS. Recent advances in pretreatment of lignocellulosic wastes and production of value added products. African Journal of Biotechnology 2009;8(8):1398-1415.
- [20] Levin DB, Islam R, Cicek N, Sparling R. Hydrogen production by *Clostridium thermocellum* 27405 from cellulosic biomass substrates. International Journal of Hydrogen Energy 2006;31(11):1496-1503
- [21] Burchhardt G, Ingram LO. Conversion of xylan to ethanol by ethanologenic strains of *Escherichia coli* and *Klebsiella oxytoca*. Applied and Environmental Microbiology 1992;58(4):1128-1133.
- [22] Galinier A1, Deutscher J, Martin-Verstraete I. Phosphorylation of either *crh* or *HPr* mediates binding of *CcpA* to the *Bacillus subtilis* *xyn cre* and catabolite repression of the *xyn* operon. Joernal of Molecular Biology 1999;286(2):307-314.
- [23] Bertram R, Wünsche A, Sprehe M, Hillen W. Regulated expression of *HPrK/P* does not affect carbon catabolite repression of the *xyn* operon and of *rocG* in *Bacillus subtilis*. FEMS Microbiology Letters 2006;259(1):147-152.
- [24] Roncero M I. Genes controlling xylan utilization by *Bacillus subtilis*. Joernal of Bacteriology 1983;156(1):257-263.
- [25] Chow V, Nong G, Preston JF. Structure, function, and regulation of the aldouronate utilization gene cluster from *Paenibacillus* sp. strain JDR-2. Joernal Bacteriology 2007;189(24):8863-8870.
- [26] Moriyama H, Fukusaki H, Crespo JC, Shinmyo A, Okada H. Structure and expression of genes coding for xylan-degrading enzymes of *Bacillus pumilus*. FEBS Journal 1987;166(3):539-545.

- [27] Zeigler DR. The genome sequence of *Bacillus subtilis* subsp. *spizizenii* W23: Insights into speciation within the *B. subtilis* complex and into the history of *B. subtilis* genetics. *Microbiology*. 2011;157(7):2033–2041.
- [28] Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden T. Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics* 2012;13:134. p.11.
- [29] Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3 - new capabilities and interfaces. *Nucleic Acids Research* 2012;40(15):e115
- [30] Koressaar T, Remm M. Enhancements and modifications of primer design program Primer3. *Bioinformatics* 2007;23(10):1289–1291
- [31] Fürste JP, Pansegrau W, Frank R, Blöcker H, Scholz P, Bagdasarian M, et al. Molecular cloning of the plasmid RP4 primase region in a multi-host-range *tacP* expression vector. *Gene*. 1986;48(1):119–131.
- [32] Sambrook J, Fritsch EF, Maniatis T, *Molecular cloning: A laboratory manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 2008.
- [33] Wahjudi M, Papaioannou E, Hendrawati O, van Assen AH, van Merkerk R, Cool RH, et al. PA0305 of *Pseudomonas aeruginosa* is a quorum quenching acylhomoserine lactone acylase belonging to the Ntn hydrolase superfamily. *Microbiology* 2011;157(Pt7):2042–2055.
- [34] Teather RM, Wood PJ. Use of Congo red polysaccharide interactions in enumeration and characterization of cellulolytic bacteria from the bovine rumen. *Applied Environmental Microbiology* 1982;43(4):777–780.
- [35] Uwe H, Chris S. A sequence property approach to searching protein databases. *Journal of Molecular Biology*. 1995;251(3):390–399.
- [36] Hiller K, Grote A, Scheer M, Münch R, Jahn D. PrediSi: Prediction of signal peptides and their cleavage positions. *Nucleic Acids Research*. 2004;32(Web Server issue):W375–W379.
- [37] Yu CS, Chen YC, Lu CH, Hwang JK: Prediction of protein subcellular localization. *Proteins: Structure, Function and Bioinformatics* 2006, 64(3):643–651.
- [38] CBS, Center for Biological Sequence Analysis. TMHMM server v. 2.0. Prediction of transmembrane helices in proteins. [Online] from <http://www.cbs.dtu.dk/services/TMHMM-2.0/> (2017) [Accessed on Dec. 25th 2016]
- [39] Softberry. Protein structure and functions. [Online] from www.softberry.com (2017) [Accessed on Dec. 25th 2016]
- [40] Zhang W, Yang M, Yang Y, Zhan J, Zhou Y, Zhao X. Optimal secretion of alkali-tolerant xylanase in *Bacillus subtilis* by signal peptide screening. *Applied Microbiology and Biotechnology* 2016;100(20):8745–8756.